

# POLYNOMIAL EQUATIONS AND RANK OF MATRICES OVER $\mathbb{F}_2$ RELATED TO PERSYMMETRIC MATRICES

JORGEN CHERLY

RÉSUMÉ. Dans cet article nous illustrons par quelques exemples la connexion entre le nombre de solutions d'équations polynômiales satisfaisant des conditions de degrés et le nombre de rang  $i$  matrices rattachées aux matrices persymétriques

ABSTRACT. In this paper we illustrate by some examples the connection between the number of solutions of polynomial equations satisfying degree conditions and the number of rank  $i$  matrices related to persymmetric matrices.

## CONTENTS

|                                                                            |    |
|----------------------------------------------------------------------------|----|
| 1. Notations                                                               | 3  |
| 2. Polynomial equations related to persymmetric matrices                   | 5  |
| 3. Polynomial equations related to double persymmetric matrices            | 8  |
| 4. Polynomial equations related to triple persymmetric matrices            | 16 |
| 5. The inverse problem                                                     | 21 |
| 5.1. The inverse problem in the case $s = 2$ , $m = 3$ , $l \geq 4$        | 22 |
| 5.2. The inverse problem in the case $s = 2$ , $m = 3$ , $l = 4$ , $k = 6$ | 24 |
| 5.3. The inverse problem in the case $s = 3$ , $m = 0$ , $l \geq 0$        | 25 |
| References                                                                 | 32 |

## 1. NOTATIONS

A matrix  $[\alpha_{i,j}]$  is persymmetric if  $\alpha_{i,j} = \alpha_{r,s}$  for  $i+j=r+s$ .

We denote by  $\Gamma_i^{s \times k}$  the number of rank  $i$  persymmetric  $s \times k$  matrices over  $\mathbb{F}_2$  of the form :

$$\begin{pmatrix} \alpha_1 & \alpha_2 & \alpha_3 & \dots & \alpha_{k-1} & \alpha_k \\ \alpha_2 & \alpha_3 & \alpha_4 & \dots & \alpha_k & \alpha_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \alpha_{s-1} & \alpha_s & \alpha_{s+1} & \dots & \alpha_{k+s-3} & \alpha_{k+s-2} \\ \alpha_s & \alpha_{s+1} & \alpha_{s+2} & \dots & \alpha_{k+s-2} & \alpha_{k+s-1} \end{pmatrix}.$$

We denote by  $\Gamma_i^{\begin{bmatrix} s \\ s+m \end{bmatrix} \times k}$  the number of rank  $i$  double persymmetric  $(2s + m) \times k$  matrices over  $\mathbb{F}_2$  of the form:

$$\left( \begin{array}{cccccc} \alpha_1 & \alpha_2 & \alpha_3 & \dots & \alpha_{k-1} & \alpha_k \\ \alpha_2 & \alpha_3 & \alpha_4 & \dots & \alpha_k & \alpha_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \alpha_{s-1} & \alpha_s & \alpha_{s+1} & \dots & \alpha_{s+k-3} & \alpha_{s+k-2} \\ \alpha_s & \alpha_{s+1} & \alpha_{s+2} & \dots & \alpha_{s+k-2} & \alpha_{s+k-1} \\ \hline \beta_1 & \beta_2 & \beta_3 & \dots & \beta_{k-1} & \beta_k \\ \beta_2 & \beta_3 & \beta_4 & \dots & \beta_k & \beta_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{m+1} & \beta_{m+2} & \beta_{m+3} & \dots & \beta_{k+m-1} & \beta_{k+m} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{s+m-1} & \beta_{s+m} & \beta_{s+m+1} & \dots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\ \beta_{s+m} & \beta_{s+m+1} & \beta_{s+m+2} & \dots & \beta_{s+m+k-2} & \beta_{s+m+k-1} \end{array} \right).$$

We denote by  $\Gamma_i^{\begin{bmatrix} s \\ s+m+l \end{bmatrix} \times k}$  the number of rank  $i$  double persymmetric  $(3s + 2m + l) \times k$  matrices over  $\mathbb{F}_2$  of the form:

$$\begin{pmatrix}
\alpha_1 & \alpha_2 & \cdots & \alpha_{k-1} & \alpha_k \\
\alpha_2 & \alpha_3 & \cdots & \alpha_k & \alpha_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots \\
\alpha_{s-1} & \alpha_s & \cdots & \alpha_{s+k-3} & \alpha_{s+k-2} \\
\alpha_s & \alpha_{s+1} & \cdots & \alpha_{s+k-2} & \alpha_{s+k-1} \\
\hline
\beta_1 & \beta_2 & \cdots & \beta_{k-1} & \beta_k \\
\beta_2 & \beta_3 & \cdots & \beta_k & \beta_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{m+1} & \beta_{m+2} & \cdots & \beta_{k+m-1} & \beta_{k+m} \\
\vdots & \vdots & \vdots & \vdots & \vdots \\
\beta_{s+m-1} & \beta_{s+m} & \cdots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\
\beta_{s+m} & \beta_{s+m+1} & \cdots & \beta_{s+m+k-2} & \beta_{s+m+k-1} \\
\hline
\gamma_1 & \gamma_2 & \cdots & \gamma_{k-1} & \gamma_k \\
\gamma_2 & \gamma_3 & \cdots & \gamma_k & \gamma_{k+1} \\
\vdots & \vdots & \vdots & \vdots & \vdots \\
\gamma_{m+1} & \gamma_{m+2} & \cdots & \gamma_{k+m-1} & \gamma_{k+m} \\
\vdots & \vdots & \vdots & \vdots & \vdots \\
\gamma_{s+m-1} & \gamma_{s+m} & \cdots & \gamma_{s+m+k-3} & \gamma_{s+m+k-2} \\
\gamma_{s+m} & \gamma_{s+m+1} & \cdots & \gamma_{s+m+k-2} & \gamma_{s+m+k-1} \\
\gamma_{s+m+1} & \gamma_{s+m+2} & \cdots & \gamma_{s+m+k-1} & \gamma_{s+m+k} \\
\vdots & \vdots & \vdots & \vdots & \vdots \\
\gamma_{s+m+l} & \gamma_{s+m+l+1} & \cdots & \gamma_{s+m+l+k-2} & \gamma_{s+m+l+k-1}
\end{pmatrix}.$$

## 2. POLYNOMIAL EQUATIONS RELATED TO PERSYMMETRIC MATRICES

*Example.* The number  $\Gamma_i^{5 \times 5}$  of persymmetric  $5 \times 5$  matrices over  $\mathbb{F}_2$  of rank  $i$  of the form

$$\begin{pmatrix} \alpha_1 & \alpha_2 & \alpha_3 & \alpha_4 & \alpha_5 \\ \alpha_2 & \alpha_3 & \alpha_4 & \alpha_5 & \alpha_6 \\ \alpha_3 & \alpha_4 & \alpha_5 & \alpha_6 & \alpha_7 \\ \alpha_4 & \alpha_5 & \alpha_6 & \alpha_7 & \alpha_8 \\ \alpha_5 & \alpha_6 & \alpha_7 & \alpha_8 & \alpha_9 \end{pmatrix}.$$

is equal to

$$\begin{cases} 1 & \text{if } i = 0, \\ 3 & \text{if } i = 1, \\ 12 & \text{if } i = 2, \\ 48 & \text{if } i = 3, \\ 192 & \text{if } i = 4, \\ 256 & \text{if } i = 5. \end{cases}$$

The general formula is given by D.E. Daykin [1] and in [2] in the case  $q = 2$ .

**Application:** The number  $R_q$  of solutions  $(Y_1, Z_1, \dots, Y_q, Z_q) \in (\mathbb{F}_2[T])^{2q}$  of the polynomial equation

$$Y_1 Z_1 + Y_2 Z_2 + \dots + Y_q Z_q = 0$$

satisfying the degree conditions

$$\deg Y_j \leq 4, \quad \deg Z_j \leq 4 \quad \text{for } 1 \leq j \leq q,$$

are connected with the numbers  $\Gamma_i$  of persymmetric  $5 \times 5$  matrices over  $\mathbb{F}_2$  of rank  $i$ , in the way that the number  $R_q$  can be written as a linear combination of the  $\Gamma_i$ .

More precisely:

$$R_q = 2^{10q-9} \sum_{i=0}^5 \Gamma_i 2^{-qi} = \begin{cases} 63 & \text{if } q = 1, \\ 8704 & \text{if } q = 2, \\ 2^{10q-9} [1 + 3 \cdot 2^{-q} + 12 \cdot 2^{-2q} + 48 \cdot 2^{-3q} + 192 \cdot 2^{-4q} + 256 \cdot 2^{-5q}] & \text{if } 3 \leq q. \end{cases}$$

*Example.* The number  $\Gamma_i = \Gamma_i^{2 \times 2}$  of persymmetric  $2 \times 2$  matrices over  $\mathbb{F}_2$  of rank  $i$  of the form

$$\begin{pmatrix} \alpha_1 & \alpha_2 \\ \alpha_2 & \alpha_3 \end{pmatrix}.$$

is equal to

$$\begin{cases} 1 & \text{if } i = 0, \\ 3 & \text{if } i = 1, \\ 4 & \text{if } i = 2, \end{cases}$$

**Application:** The number  $R_q$  of solutions  $(Y_1, Z_1, \dots, Y_q, Z_q) \in (\mathbb{F}_2[T])^{2q}$  of the polynomial equation

$$Y_1 Z_1 + Y_2 Z_2 + \dots + Y_q Z_q = 0$$

satisfying the degree conditions

$$\deg Y_j \leq 1, \quad \deg Z_j \leq 1 \quad \text{for } 1 \leq j \leq q,$$

is connected with the numbers  $\Gamma_i$  of persymmetric  $2 \times 2$  matrices over  $\mathbb{F}_2$  of rank  $i$ , in the way that the number  $R_q$  can be written as a linear combination of the  $\Gamma_i$ .

More precisely:

$$\begin{aligned} R_q &= 2^{4q-3} \sum_{i=0}^2 \Gamma_i 2^{-qi} \\ &= \begin{cases} 7 & \text{if } q = 1, \\ 64 & \text{if } q = 2, \\ 2^{4q-3} [1 + 3 \cdot 2^{-q} + 4 \cdot 2^{-2q}] & \text{if } 3 \leq q. \end{cases} \end{aligned}$$

*Example.* The fraction of square persymmetric matrices which are invertible is equal to  $\frac{1}{2}$

*Example.* The number  $\Gamma_i^{\begin{bmatrix} 5 \\ 1 \end{bmatrix} \times 5}$  of rank  $i$  matrices of the form

$$\begin{pmatrix} \alpha_1 & \alpha_2 & \alpha_3 & \alpha_4 & \alpha_5 \\ \alpha_2 & \alpha_3 & \alpha_4 & \alpha_5 & \alpha_6 \\ \alpha_3 & \alpha_4 & \alpha_5 & \alpha_6 & \alpha_7 \\ \alpha_4 & \alpha_5 & \alpha_6 & \alpha_7 & \alpha_8 \\ \alpha_5 & \alpha_6 & \alpha_7 & \alpha_8 & \alpha_9 \\ \hline \beta_1 & \beta_2 & \beta_3 & \beta_4 & \beta_5 \end{pmatrix}$$

is equal to

$$2^i \Gamma_i^{5 \times 5} + (2^5 - 2^{i-1}) \cdot \Gamma_{i-1}^{5 \times 5} \quad \text{for } 0 \leq i \leq 5,$$

$$= \begin{cases} 1 & \text{if } i = 0, \\ 37 & \text{if } i = 1, \\ 138 & \text{if } i = 2, \\ 720 & \text{if } i = 3, \\ 4224 & \text{if } i = 4, \\ 11264 & \text{if } i = 5. \end{cases}$$

**Application:** The number  $R_q$  of solutions  $(Y_1, Z_1, U_1, Y_2, Z_2, U_2, \dots, Y_q, Z_q, U_q) \in (\mathbb{F}_2[T])^{3q}$

of the polynomial equations

$$\begin{cases} Y_1 Z_1 + Y_2 Z_2 + \dots + Y_q Z_q = 0 \\ Y_1 U_1 + Y_2 U_2 + \dots + Y_q U_q = 0 \end{cases}$$

satisfying the degree conditions

$$\deg Y_j \leq 4, \quad \deg Z_j \leq 4, \quad \deg U_j \leq 0, \quad \text{for } 1 \leq j \leq q$$

is connected with the numbers  $\Gamma_i^{\begin{bmatrix} 5 \\ 1 \end{bmatrix} \times 5}$ , in the way that the number  $R_q$  can

be written as a linear combination of the  $\Gamma_i^{\begin{bmatrix} 5 \\ 1 \end{bmatrix} \times 5}$

More precisely:

$$\begin{aligned} R_q &= 2^{11q-14} \sum_{i=0}^5 \Gamma_i^{\begin{bmatrix} 5 \\ 1 \end{bmatrix} \times 5} 2^{-iq} \\ &= \begin{cases} 95 & \text{if } q = 1, \\ 14752 & \text{if } q = 2, \\ 2^{11q-14} [1 + 37 \cdot 2^{-q} + 138 \cdot 2^{-2q} + 720 \cdot 2^{-3q} + 4224 \cdot 2^{-4q} + 11264 \cdot 2^{-5q}] & \text{if } 3 \leq q. \end{cases} \end{aligned}$$

*Example.* The number  $\Gamma_i^{\left[\begin{smallmatrix} 2 \\ (2) \end{smallmatrix}\right] \times 4}$  of rank  $i$  matrices of the form

$$\begin{pmatrix} \alpha_1 & \alpha_2 & \alpha_3 & \alpha_4 \\ \alpha_4 & \alpha_5 & \alpha_6 & \alpha_7 \\ \alpha_7 & \alpha_8 & \alpha_9 & \alpha_{10} \\ \alpha_{10} & \alpha_{11} & \alpha_{12} & \alpha_{13} \end{pmatrix} \underset{\text{rank}}{\sim} \begin{pmatrix} \alpha_1 & \alpha_2 & \alpha_3 & \alpha_4 \\ \alpha_2 & \alpha_3 & \alpha_4 & \alpha_5 \\ \beta_{11} & \beta_{12} & \beta_{13} & \beta_{14} \\ \beta_{21} & \beta_{22} & \beta_{23} & \beta_{24} \end{pmatrix}$$

is equal to

$$2^{2i} \cdot \Gamma_i^{2 \times 4} + 3 \cdot 2^{i-1} \cdot (2^4 - 2^{i-1}) \cdot \Gamma_{i-1}^{2 \times 4} + (2^4 - 2^{i-1}) \cdot (2^4 - 2^{i-2}) \cdot \Gamma_{i-2}^{2 \times 4} \quad \text{for } 0 \leq i \leq 4$$

$$= \begin{cases} 1 & \text{if } i = 0, \\ 57 & \text{if } i = 1, \\ 910 & \text{if } i = 2, \\ 4536 & \text{if } i = 3, \\ 2688 & \text{if } i = 4. \end{cases}$$

**Application:** The number  $R_q$  of solutions  $(Y_1, Z_1, U_1, V_1, Y_2, Z_2, U_2, V_2, \dots, Y_q, Z_q, U_q, V_q) \in (\mathbb{F}_2[T])^{4q}$

of the polynomial equations

$$\begin{cases} Y_1 Z_1 + Y_2 Z_2 + \dots + Y_q Z_q = 0 \\ Y_1 U_1 + Y_2 U_2 + \dots + Y_q U_q = 0 \\ Y_1 V_1 + Y_2 V_2 + \dots + Y_q V_q = 0 \end{cases}$$

satisfying the degree conditions

$$\deg Y_j \leq 3, \quad \deg Z_j \leq 1, \quad \deg U_j \leq 0, \quad \deg V_j \leq 0 \quad \text{for } 1 \leq j \leq q$$

is equal to

$$2^{8q-13} \sum_{i=0}^4 \Gamma_i^{\left[\begin{smallmatrix} 2 \\ (2) \end{smallmatrix}\right] \times 4} \cdot 2^{-iq} = 2^{8q-13} [1 + 57 \cdot 2^{-q} + 910 \cdot 2^{-2q} + 4536 \cdot 2^{-3q} + 2688 \cdot 2^{-4q}]$$

For more details see [2]

### 3. POLYNOMIAL EQUATIONS RELATED TO DOUBLE PERSYMMETRIC MATRICES

*Example.* The most simple problem concerning double persymmetric matrices with entries in  $\mathbb{F}_2$ :

Compute the number  $\Gamma_i^{\left[\begin{smallmatrix} 2 \\ 2 \end{smallmatrix}\right] \times k}$  of double persymmetric matrices in  $\mathbb{F}_2$  of rank



$0 \leq i \leq \inf(4, k)$  of the form:

$$\begin{pmatrix} \alpha_1 & \alpha_2 & \dots & \alpha_{k-1} & \alpha_k \\ \alpha_2 & \alpha_3 & \dots & \alpha_k & \alpha_{k+1} \\ \hline \beta_1 & \beta_2 & \dots & \beta_{k-1} & \beta_k \\ \beta_2 & \beta_3 & \dots & \beta_k & \beta_{k+1} \end{pmatrix}.$$

We get

$$\Gamma_i^{\left[\begin{smallmatrix} 2 \\ 2 \end{smallmatrix}\right] \times k} = \begin{cases} 1 & \text{if } i = 0, k \geq 1, \\ 9 & \text{if } i = 1, k > 1, \\ 3 \cdot 2^{k+1} + 30 & \text{if } i = 2, k > 2, \\ 21 \cdot 2^{k+1} - 168 & \text{if } i = 3, k > 3, \\ 2^{2k+2} - 3 \cdot 2^{k+4} + 128 & \text{if } i = 4, k \geq 4 \end{cases}$$

$$\Gamma_i^{\left[\begin{smallmatrix} 2 \\ 2 \end{smallmatrix}\right] \times i} = \begin{cases} 15 & \text{if } i = 1, \\ 54 & \text{if } i = 2, \\ 168 & \text{if } i = 3, \\ 384 & \text{if } i = 4 \end{cases}$$

**Application:** The number  $R_q$  of solutions  $(Y_1, Z_1, U_1, Y_2, Z_2, U_2, \dots, Y_q, Z_q, U_q) \in (\mathbb{F}_2[T])^{3q}$

of the polynomial equations

$$\begin{cases} Y_1 Z_1 + Y_2 Z_2 + \dots + Y_q Z_q = 0 \\ Y_1 U_1 + Y_2 U_2 + \dots + Y_q U_q = 0 \end{cases}$$

satisfying the degree conditions

$$\deg Y_j \leq k-1, \quad \deg Z_j \leq 1, \quad \deg U_j \leq 1, \quad \text{for } 1 \leq j \leq q$$

is connected with the numbers  $\Gamma_i^{\left[\begin{smallmatrix} 2 \\ 2 \end{smallmatrix}\right] \times k}$ , in the way that the number  $R_q$  can

be written as a linear combination of the  $\Gamma_i^{\left[\begin{smallmatrix} 2 \\ 2 \end{smallmatrix}\right] \times k}$

More precisely:

$$\begin{aligned} R_q &= 2^{(k+4)q-2k-2} \sum_{i=0}^{\inf(4,k)} \Gamma_i^{\left[\begin{smallmatrix} 2 \\ 2 \end{smallmatrix}\right] \times k} 2^{-iq} \\ &= \begin{cases} 2^k + 15 & \text{if } q = 1, k \geq 4 \\ 2^{2k+8} + 27 \cdot 2^{k+1} + 192 & \text{if } q = 2, k \geq 4 \\ 2^{kq+4q-2k-2} \cdot [1 + 9 \cdot 2^{-q} + (3 \cdot 2^{k+1} + 30) \cdot 2^{-2q} + (21 \cdot 2^{k+1} - 168) \cdot 2^{-3q}] & \text{if } 3 \leq q, k \geq 4 \\ + 2^{kq+4q-2k-2} \cdot [(2^{2k+2} - 3 \cdot 2^{k+4} + 128) \cdot 2^{-4q}] & \end{cases} \end{aligned}$$

**The case  $k = 1$**

$$R_q = 2^{5q-4} \sum_{i=0}^1 \Gamma_i^{\left[\begin{smallmatrix} 2 \\ 2 \end{smallmatrix}\right] \times 1} 2^{-iq} = 2^{5q-4} + 15 \cdot 2^{4q-4}$$

**The case  $k = 2$**

$$R_q = 2^{6q-6} \sum_{i=0}^2 \Gamma_i^{\left[\begin{smallmatrix} 2 \\ 2 \end{smallmatrix}\right] \times 2} 2^{-iq} = 2^{6q-6} + 9 \cdot 2^{5q-6} + 54 \cdot 2^{4q-6}$$

**The case  $k = 3$**

$$R_q = 2^{7q-8} \sum_{i=0}^3 \Gamma_i^{\left[\begin{smallmatrix} 2 \\ 2 \end{smallmatrix}\right] \times 3} 2^{-iq} = 2^{7q-8} + 9 \cdot 2^{6q-8} + 78 \cdot 2^{5q-8} + 168 \cdot 2^{4q-8}$$

*Example.* The number  $\Gamma_i^{\left[\begin{smallmatrix} 5 \\ 5 \end{smallmatrix}\right] \times k}$  of double persymmetric  $10 \times k$  matrices over  $\mathbb{F}_2$  of rank  $i$  of the form

$$\left( \begin{array}{ccccc} \alpha_1 & \alpha_2 & \dots & \alpha_{k-1} & \alpha_k \\ \alpha_2 & \alpha_3 & \dots & \alpha_k & \alpha_{k+1} \\ \alpha_3 & \alpha_4 & \dots & \alpha_{k+1} & \alpha_{k+2} \\ \alpha_4 & \alpha_5 & \dots & \alpha_{k+2} & \alpha_{k+3} \\ \alpha_5 & \alpha_6 & \dots & \alpha_{k+3} & \alpha_{k+4} \\ \hline \beta_1 & \beta_2 & \dots & \beta_{k-1} & \beta_k \\ \beta_2 & \beta_3 & \dots & \beta_k & \beta_{k+1} \\ \beta_3 & \beta_4 & \dots & \beta_{k+1} & \beta_{k+2} \\ \beta_4 & \beta_5 & \dots & \beta_{k+2} & \beta_{k+3} \\ \beta_5 & \beta_6 & \dots & \beta_{k+3} & \beta_{k+4} \end{array} \right).$$

is equal to

$$\left\{ \begin{array}{ll} 1 & \text{if } i = 0, k \geq 1, \\ 9 & \text{if } i = 1, k > 1, \\ 78 & \text{if } i = 2, k > 2, \\ 648 & \text{if } i = 3, k > 3, \\ 5280 & \text{if } i = 4, k > 4, \\ 3 \cdot 2^{k+4} + 39552 & \text{if } i = 5, k > 5, \\ 21 \cdot 2^{k+4} + 290304 & \text{if } i = 6, k > 6, \\ 21 \cdot 2^{k+7} + 1892352 & \text{if } i = 7, k > 7, \\ 21 \cdot 2^{k+10} + 8257536 & \text{if } i = 8, k > 8, \\ 21 \cdot 2^{k+13} - 44040192 & \text{if } i = 9, k > 9, \\ 2^{2k+8} - 3 \cdot 2^{k+16} + 2^{25} & \text{if } i = 10, k \geq 10. \end{array} \right.$$

**The case  $k=i$**

$$\Gamma_i^{\begin{bmatrix} 5 \\ 5 \end{bmatrix} \times i} = \begin{cases} 2^{10} - 1 & \text{if } i = 1, \\ 2^{12} - 10 & \text{if } i = 2, \\ 2^{14} - 88 & \text{if } i = 3, \\ 2^{16} - 736 & \text{if } i = 4, \\ 2^{18} - 6016 & \text{if } i = 5, \\ 2^{20} - 48640 & \text{if } i = 6, \\ 2^{22} - 385024 & \text{if } i = 7, \\ 2^{24} - 3014656 & \text{if } i = 8, \\ 2^{26} - 23068672 & \text{if } i = 9, \\ 2^{28} - 167772160 & \text{if } i = 10, \end{cases}$$

**The case  $k=4$**

The number  $\Gamma_i^{\begin{bmatrix} 5 \\ 5 \end{bmatrix} \times 4}$  of double persymmetric  $10 \times 4$  matrices over  $\mathbb{F}_2$  of rank  $i$  of the form

$$\begin{pmatrix} \alpha_1 & \alpha_2 & \alpha_3 & \alpha_4 \\ \alpha_2 & \alpha_3 & \alpha_4 & \alpha_5 \\ \alpha_3 & \alpha_4 & \alpha_5 & \alpha_6 \\ \alpha_4 & \alpha_5 & \alpha_6 & \alpha_7 \\ \alpha_5 & \alpha_6 & \alpha_7 & \alpha_8 \\ \hline \beta_1 & \beta_2 & \beta_3 & \beta_4 \\ \beta_2 & \beta_3 & \beta_4 & \beta_5 \\ \beta_3 & \beta_4 & \beta_5 & \beta_6 \\ \beta_4 & \beta_5 & \beta_6 & \beta_7 \\ \beta_5 & \beta_6 & \beta_7 & \beta_8 \end{pmatrix}.$$

is equal to

$$\begin{cases} 1 & \text{if } i = 0, \\ 9 & \text{if } i = 1 \\ 78 & \text{if } i = 2, \\ 648 & \text{if } i = 3 \\ 64800 & \text{if } i = 4 \end{cases}$$

**Application:** The number  $R_q$  of solutions  $(Y_1, Z_1, U_1, Y_2, Z_2, U_2, \dots, Y_q, Z_q, U_q) \in (\mathbb{F}_2[T])^{3q}$

of the polynomial equations

$$\begin{cases} Y_1 Z_1 + Y_2 Z_2 + \dots + Y_q Z_q = 0 \\ Y_1 U_1 + Y_2 U_2 + \dots + Y_q U_q = 0 \end{cases}$$

satisfying the degree conditions

$$\deg Y_j \leq 3, \quad \deg Z_j \leq 4, \quad \deg U_j \leq 4, \quad \text{for } 1 \leq j \leq q$$

is connected with the numbers  $\Gamma_i^{\begin{bmatrix} 5 \\ 5 \end{bmatrix} \times 4}$ , in the way that the number  $R_q$  can be written as a linear combination of the  $\Gamma_i^{\begin{bmatrix} 5 \\ 5 \end{bmatrix} \times 4}$ . More precisely:

$$\begin{aligned} R_q &= 2^{14q-16} \sum_{i=0}^4 \Gamma_i^{\begin{bmatrix} 5 \\ 5 \end{bmatrix} \times 4} 2^{-iq} \\ &= 2^{14q-16} \cdot [1 + 9 \cdot 2^{-q} + 78 \cdot 2^{-2q} + 648 \cdot 2^{-3q} + 64800 \cdot 2^{-4q}] \end{aligned}$$

*Example.* The fraction of square double persymmetric matrices which are invertible is equal to  $\frac{3}{8}$

*Example.* The number  $\Gamma_i^{\begin{bmatrix} 2 & 2 \\ 2+3 \end{bmatrix} \times 4}$  of double persymmetric  $7 \times 4$  matrices over  $\mathbb{F}_2$  of rank  $i$  of the form

$$\begin{pmatrix} \alpha_1 & \alpha_2 & \alpha_3 & \alpha_4 \\ \alpha_2 & \alpha_3 & \alpha_4 & \alpha_5 \\ \hline \beta_1 & \beta_2 & \beta_3 & \beta_4 \\ \beta_2 & \beta_3 & \beta_4 & \beta_5 \\ \beta_3 & \beta_4 & \beta_5 & \beta_6 \\ \beta_4 & \beta_5 & \beta_6 & \beta_7 \\ \beta_5 & \beta_6 & \beta_7 & \beta_8 \end{pmatrix}.$$

is given by

$$\begin{cases} 1 & \text{if } i = 0, \\ 9 & \text{if } i = 1, \\ 94 & \text{if } i = 2, \\ 600 & \text{if } i = 3, \\ 7488 & \text{if } i = 4 \end{cases}$$

### Application:

The number  $R_q$  of solutions  $(Y_1, Z_1, U_1, Y_2, Z_2, U_2, \dots, Y_q, Z_q, U_q) \in (\mathbb{F}_2[T])^{3q}$

of the polynomial equations

$$\begin{cases} Y_1 Z_1 + Y_2 Z_2 + \dots + Y_q Z_q = 0 \\ Y_1 U_1 + Y_2 U_2 + \dots + Y_q U_q = 0 \end{cases}$$

satisfying the degree conditions

$$\deg Y_j \leq 3, \quad \deg Z_j \leq 1, \quad \deg U_j \leq 4, \quad \text{for } 1 \leq j \leq q$$

is connected with the numbers  $\Gamma_i^{\left[\begin{smallmatrix} 2 & 2 \\ 2+3 \end{smallmatrix}\right] \times 4}$ , in the way that the number  $R_q$  can be written as a linear combination of the  $\Gamma_i^{\left[\begin{smallmatrix} 2 & 2 \\ 2+3 \end{smallmatrix}\right] \times 4}$ . More precisely:

$$\begin{aligned} R_q &= 2^{11q-13} \sum_{i=0}^4 \Gamma_i^{\left[\begin{smallmatrix} 2 & 2 \\ 2+3 \end{smallmatrix}\right] \times 4} 2^{-iq} \\ &= 2^{11q-13} \cdot [1 + 9 \cdot 2^{-q} + 94 \cdot 2^{-2q} + 600 \cdot 2^{-3q} + 7488 \cdot 2^{-4q}] \end{aligned}$$

*Example.* The number  $\Gamma_i^{\left[\begin{smallmatrix} 2 & 2 \\ 2+3 \\ (1) \end{smallmatrix}\right] \times 4}$  of rank  $i$   $8 \times 4$  matrices over  $\mathbb{F}_2$  of the form

$$\begin{pmatrix} \alpha_1 & \alpha_2 & \alpha_3 & \alpha_4 \\ \alpha_2 & \alpha_3 & \alpha_4 & \alpha_5 \\ \hline \beta_1 & \beta_2 & \beta_3 & \beta_4 \\ \beta_2 & \beta_3 & \beta_4 & \beta_5 \\ \beta_3 & \beta_4 & \beta_5 & \beta_6 \\ \beta_4 & \beta_5 & \beta_6 & \beta_7 \\ \beta_5 & \beta_6 & \beta_7 & \beta_8 \\ \hline \gamma_{11} & \gamma_{12} & \gamma_{13} & \gamma_4 \end{pmatrix}.$$

is equal to

$$\begin{aligned} &2^i \cdot \Gamma_i^{\left[\begin{smallmatrix} 2 & 2 \\ 2+3 \end{smallmatrix}\right] \times 4} + (2^4 - 2^{i-1}) \cdot \Gamma_{i-1}^{\left[\begin{smallmatrix} 2 & 2 \\ 2+3 \end{smallmatrix}\right] \times 4} \quad \text{for } 1 \leq i \leq 4 \\ &= \begin{cases} 1 & \text{if } i = 0, \\ 33 & \text{if } i = 1, \\ 502 & \text{if } i = 2, \\ 5928 & \text{if } i = 3, \\ 124608 & \text{if } i = 4 \end{cases} \end{aligned}$$

### Application:

The number  $R_q$  of solutions  $(Y_1, Z_1, U_1, V_1, \dots, Y_q, Z_q, U_q, V_q) \in (\mathbb{F}_2[T])^{4q}$  of the polynomial equations

$$\begin{cases} Y_1 Z_1 + Y_2 Z_2 + \dots + Y_q Z_q = 0, \\ Y_1 U_1 + Y_2 U_2 + \dots + Y_q U_q = 0, \\ Y_1 V_1 + Y_2 V_2 + \dots + Y_q V_q = 0, \end{cases}$$

satisfying the degree conditions

$$\deg Y_j \leq 3, \quad \deg Z_j \leq 1, \quad \deg U_j \leq 4, \quad \deg V_j \leq 0 \quad \text{for } 1 \leq j \leq q$$

is connected with the numbers  $\Gamma_i^{\left[ \begin{smallmatrix} 2 \\ 2+3 \\ (1) \end{smallmatrix} \right] \times 4}$ , in the way that the number  $R_q$  can be written as a linear combination of the  $\Gamma_i^{\left[ \begin{smallmatrix} 2 \\ 2+3 \\ (1) \end{smallmatrix} \right] \times 4}$

More precisely:

$$\begin{aligned} R_q &= 2^{12q-17} \sum_{i=0}^4 \Gamma_i^{\left[ \begin{smallmatrix} 2 \\ 2+3 \\ (1) \end{smallmatrix} \right] \times 4} 2^{-iq} \\ &= 2^{12q-17} \cdot [1 + 33 \cdot 2^{-q} + 502 \cdot 2^{-2q} + 5928 \cdot 2^{-3q} + 124608 \cdot 2^{-4q}] \end{aligned}$$

*Example.* The number  $\Gamma_i^{\left[ \begin{smallmatrix} 2 \\ 2 \\ (4) \end{smallmatrix} \right] \times 4}$  of rank  $i$  matrices of the form

$$\begin{pmatrix} \alpha_1 & \alpha_2 & \alpha_3 & \alpha_4 & \alpha_5 & \alpha_6 & \alpha_7 & \alpha_8 \\ \alpha_7 & \alpha_8 & \alpha_9 & \alpha_{10} & \alpha_{11} & \alpha_{12} & \alpha_{13} & \alpha_{14} \\ \alpha_{13} & \alpha_{14} & \alpha_{15} & \alpha_{16} & \alpha_{17} & \alpha_{18} & \alpha_{19} & \alpha_{20} \\ \alpha_{19} & \alpha_{20} & \alpha_{21} & \alpha_{22} & \alpha_{23} & \alpha_{24} & \alpha_{25} & \alpha_{26} \end{pmatrix} \underset{\text{rank}}{\sim} \begin{pmatrix} \alpha_1 & \alpha_2 & \alpha_3 & \alpha_4 \\ \alpha_2 & \alpha_3 & \alpha_4 & \alpha_5 \\ \hline \beta_1 & \beta_2 & \beta_3 & \beta_4 \\ \beta_2 & \beta_3 & \beta_4 & \beta_5 \\ \hline \gamma_{11} & \gamma_{12} & \gamma_{13} & \gamma_{14} \\ \gamma_{21} & \gamma_{22} & \gamma_{23} & \gamma_{24} \\ \gamma_{31} & \gamma_{32} & \gamma_{33} & \gamma_{34} \\ \gamma_{41} & \gamma_{42} & \gamma_{43} & \gamma_{44} \end{pmatrix}$$

is equal to

$$\begin{aligned} &2^{4i} \cdot \Gamma_i^{\left[ \begin{smallmatrix} 2 \\ 2 \end{smallmatrix} \right] \times 4} + 15 \cdot 2^{(i-1)3} (2^k - 2^{i-1}) \cdot \Gamma_{i-1}^{\left[ \begin{smallmatrix} 2 \\ 2 \end{smallmatrix} \right] \times 4} \\ &+ 35 \cdot 2^{2i-4} (2^k - 2^{i-1}) (2^k - 2^{i-2}) \cdot \Gamma_{i-2}^{\left[ \begin{smallmatrix} 2 \\ 2 \end{smallmatrix} \right] \times 4} \\ &+ 15 \cdot 2^{i-3} (2^k - 2^{i-1}) (2^k - 2^{i-2}) (2^k - 2^{i-3}) \Gamma_{i-3}^{\left[ \begin{smallmatrix} 2 \\ 2 \end{smallmatrix} \right] \times 4} \\ &+ (2^k - 2^{i-1}) (2^k - 2^{i-2}) (2^k - 2^{i-3}) (2^k - 2^{i-4}) \Gamma_{i-4}^{\left[ \begin{smallmatrix} 2 \\ 2 \end{smallmatrix} \right] \times 4} \text{ for } 0 \leq i \leq \inf(4, 8), \end{aligned}$$

$$= \begin{cases} 1 & \text{if } i = 0, \\ 369 & \text{if } i = 1, \\ 54726 & \text{if } i = 2, \\ 3765384 & \text{if } i = 3, \\ 63288384 & \text{if } i = 4. \end{cases}$$

Then the number of solutions  
 $(Y_1, Z_1, U_1, V_1^{(1)}, V_2^{(1)}, V_3^{(1)}, V_4^{(1)}, Y_2, Z_2, U_2, V_1^{(2)}, V_2^{(2)}, V_3^{(2)}, V_4^{(2)}, \dots, Y_q, Z_q, U_q, V_1^{(q)}, V_2^{(q)}, V_3^{(q)}, V_4^{(q)}) \in (\mathbb{F}_2[T])^{7q}$

of the polynomial equations

$$\begin{cases} Y_1 Z_1 + Y_2 Z_2 + \dots + Y_q Z_q = 0 \\ Y_1 U_1 + Y_2 U_2 + \dots + Y_q U_q = 0 \\ Y_1 V_1^{(1)} + Y_2 V_1^{(2)} + \dots + Y_q V_1^{(q)} = 0 \\ Y_1 V_2^{(1)} + Y_2 V_2^{(2)} + \dots + Y_q V_2^{(q)} = 0 \\ Y_1 V_3^{(1)} + Y_2 V_3^{(2)} + \dots + Y_q V_3^{(q)} = 0 \\ Y_1 V_4^{(1)} + Y_2 V_4^{(2)} + \dots + Y_q V_4^{(q)} = 0 \end{cases}$$

satisfying the degree conditions

$$\deg Y_i \leq 3, \quad \deg Z_i \leq 1, \quad \deg U_i \leq 1, \quad \deg V_j^i \leq 0, \quad \text{for } 1 \leq j \leq 4 \quad 1 \leq i \leq q$$

is equal to

$$2^{12q-26} \sum_{i=0}^4 \Gamma_i^{\left[ \frac{2}{(4)} \right] \times k} 2^{-iq} = 2^{12q-26} [1 + 369 \cdot 2^{-q} + 54726 \cdot 2^{-2q} + 3765384 \cdot 2^{-3q} + 63288384 \cdot 2^{-4q}]$$

For more details see [3], [4]

#### 4. POLYNOMIAL EQUATIONS RELATED TO TRIPLE PERSYMMETRIC MATRICES

*Example.* The number  $\Gamma_i^{\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \times 1}$  of triple persymmetric  $3 \times 1$  matrices over  $\mathbb{F}_2$  of rank  $i$  of the form

$$\begin{pmatrix} \frac{\alpha_1}{\beta_1} \\ \gamma_1 \end{pmatrix}.$$

is equal to

$$\begin{cases} 1 & \text{if } i = 0 \\ 7 & \text{if } i = 2 \end{cases}$$

##### **Application:**

The number of solutions  $(y_1, z_1, u_1, v_1, \dots, y_q, z_q, u_q, v_q) \in \mathbb{F}_2^{4q}$  of the following system of quadratic equations :

$$\begin{cases} y_1 z_1 + y_2 z_2 + \dots + y_q z_q = 0, \\ y_1 u_1 + y_2 u_2 + \dots + y_q u_q = 0, \\ y_1 v_1 + y_2 v_2 + \dots + y_q v_q = 0, \end{cases}$$

is equal to

$$\begin{aligned} & 2^{4q-3} \cdot \sum_{i=0}^1 \Gamma_i^{\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \times 1} \cdot 2^{-qi} \\ &= 2^{4q-3} \cdot (1 + 7 \cdot 2^{-q} +) \\ &= 2^{4q-3} + 7 \cdot 2^{3q-3} \end{aligned}$$

##### **Generalization**

The number  $\Gamma_i^{\begin{bmatrix} 1 \\ \vdots \\ 1 \end{bmatrix} \times 3}$  of  $n$ -times persymmetric  $n \times 1$  matrices over  $\mathbb{F}_2$  of rank  $i$  of the form

$$\begin{pmatrix} \frac{\alpha_1}{\alpha_2} \\ \vdots \\ \alpha_n \end{pmatrix}.$$

is equal to

$$\begin{cases} 1 & \text{if } i = 0 \\ 2^n - 1 & \text{if } i = 2 \end{cases}$$



**Application:**

The number of solutions  $(y_1, z_1^{(1)}, z_1^{(2)}, \dots, z_1^{(n)}, \dots, y_q, z_q^{(1)}, z_q^{(2)}, \dots, z_q^{(n)}) \in \mathbb{F}_2^{(n+1)q}$  of the following system of quadratic equations :

$$\begin{cases} y_1 z_1^{(1)} + y_2 z_2^{(1)} + \dots + y_q z_q^{(1)} = 0, \\ y_1 z_1^{(2)} + y_2 z_2^{(2)} + \dots + y_q z_q^{(2)} = 0, \\ y_1 z_1^{(3)} + y_2 z_2^{(3)} + \dots + y_q z_q^{(3)} = 0, \\ \vdots \\ y_1 z_1^{(n)} + y_2 z_2^{(n)} + \dots + y_q z_q^{(n)} = 0 \end{cases}$$

is equal to

$$\begin{aligned} & 2^{(n+1)q-n} \cdot \sum_{i=0}^1 \Gamma_i^{\begin{bmatrix} 1 \\ \vdots \\ i \end{bmatrix} \times 3} \cdot 2^{-qi} \\ &= 2^{(n+1)q-n} \cdot (1 + (2^n - 1) \cdot 2^{-q}) = 2^{nq-n} \cdot [2^q + 2^n - 1] \end{aligned}$$

*Example.* The number  $\Gamma_i^{\begin{bmatrix} 2 \\ 2 \end{bmatrix} \times 6}$  of triple persymmetric  $6 \times 6$  matrices over  $\mathbb{F}_2$  of rank  $i$  of the form

$$\begin{pmatrix} \alpha_1 & \alpha_2 & \alpha_3 & \alpha_4 & \alpha_5 & \alpha_6 \\ \alpha_2 & \alpha_3 & \alpha_4 & \alpha_5 & \alpha_6 & \alpha_7 \\ \hline \beta_1 & \beta_2 & \beta_3 & \beta_4 & \beta_5 & \beta_6 \\ \beta_2 & \beta_3 & \beta_3 & \beta_5 & \beta_6 & \beta_7 \\ \hline \gamma_1 & \gamma_2 & \gamma_3 & \gamma_4 & \gamma_5 & \gamma_6 \\ \gamma_2 & \gamma_3 & \gamma_4 & \gamma_5 & \gamma_6 & \gamma_7 \end{pmatrix}.$$

is equal to

$$\begin{cases} 1 & \text{if } i = 0 \\ 21 & \text{if } i = 1 \\ 1162 & \text{if } i = 2 \\ 20160 & \text{if } i = 3 \\ 258720 & \text{if } i = 4 \\ 1128960 & \text{if } i = 5 \\ 688128 & \text{if } i = 6 \end{cases}$$

**Application:**

The number  $R_q$  of solutions  $(Y_1, Z_1, U_1, V_1, \dots, Y_q, Z_q, U_q, V_q) \in (\mathbb{F}_2[T])^{4q}$  of the polynomial equations

$$\begin{cases} Y_1 Z_1 + Y_2 Z_2 + \dots + Y_q Z_q = 0, \\ Y_1 U_1 + Y_2 U_2 + \dots + Y_q U_q = 0, \\ Y_1 V_1 + Y_2 V_2 + \dots + Y_q V_q = 0, \end{cases}$$

satisfying the degree conditions

$$\deg Y_j \leq 5, \quad \deg Z_j \leq 1, \quad \deg U_j \leq 1, \quad \deg V_j \leq 1 \quad \text{for } 1 \leq j \leq q.$$

is equal to

$$\begin{aligned} & 2^{12q-21} \cdot \sum_{i=0}^6 \Gamma_i^{\left[\frac{2}{2}\right] \times 6} \cdot 2^{-qi} \\ &= 2^{12q-21} \cdot (1 + 21 \cdot 2^{-q} + 1162 \cdot 2^{-2q} + 20160 \cdot 2^{-3q} + 258720 \cdot 2^{-4q} + 1128960 \cdot 2^{-5q} + 688128 \cdot 2^{-6q}) \\ &= 2^{6q-21} \cdot (2^{6q} + 21 \cdot 2^{5q} + 1162 \cdot 2^{4q} + 20160 \cdot 2^{3q} + 258720 \cdot 2^{2q} + 1128960 \cdot 2^q + 688128) \end{aligned}$$

*Example.* The fraction of square triple persymmetric matrices which are invertible is equal to  $\frac{21}{64}$

*Example.* The number  $\Gamma_i^{\left[\frac{2}{2}\right] \times 2}$  of triple persymmetric  $6 \times 2$  matrices over  $\mathbb{F}_2$  of rank  $i$  of the form

$$\begin{pmatrix} \alpha_1 & \alpha_2 \\ \alpha_2 & \alpha_3 \\ \beta_1 & \beta_2 \\ \beta_2 & \beta_3 \\ \gamma_1 & \gamma_2 \\ \gamma_2 & \gamma_3 \end{pmatrix}.$$

is equal to

$$\begin{cases} 1 & \text{if } i = 0 \\ 21 & \text{if } i = 1 \\ 490 & \text{if } i = 2 \end{cases}$$

### Application:

The number  $R_q$  of solutions

$(Y_1, Z_1, U_1, V_1, \dots, Y_q, Z_q, U_q, V_q) \in (\mathbb{F}_2[T])^{4q}$  of the polynomial equations

$$\begin{cases} Y_1 Z_1 + Y_2 Z_2 + \dots + Y_q Z_q = 0, \\ Y_1 U_1 + Y_2 U_2 + \dots + Y_q U_q = 0, \\ Y_1 V_1 + Y_2 V_2 + \dots + Y_q V_q = 0, \end{cases}$$

satisfying the degree conditions

$$\deg Y_j \leq 1, \quad \deg Z_j \leq 1, \quad \deg U_j \leq 1, \quad \deg V_j \leq 1 \quad \text{for } 1 \leq j \leq q.$$

is equal to

$$\begin{aligned}
& 2^{8q-9} \cdot \sum_{i=0}^2 \Gamma_i^{\left[\frac{2}{2}\right] \times 2} \cdot 2^{-qi} \\
&= 2^{8q-9} \cdot (1 + 21 \cdot 2^{-q} + 490 \cdot 2^{-2q}) \\
&= 2^{8q-9} + 21 \cdot 2^{7q-9} + 490 \cdot 2^{6q-9}
\end{aligned}$$

**Application :** The number of solutions  $(x_1, x_2, \dots, x_{15}, x_{16}) \in \mathbb{F}_2^{16}$  of the following system of quadratic equations :

$$\left\{ \begin{array}{l} x_1x_5 + x_3x_7 = 0, \\ x_1x_9 + x_3x_{11} = 0, \\ x_1x_{13} + x_3x_{15} = 0, \\ x_1x_6 + x_2x_5 + x_3x_8 + x_4x_7 = 0 \\ x_1x_{10} + x_2x_9 + x_3x_{12} + x_4x_{11} = 0 \\ x_1x_{14} + x_2x_{13} + x_3x_{16} + x_4x_{15} = 0 \\ x_2x_6 + x_4x_8 = 0, \\ x_2x_{10} + x_4x_{12} = 0, \\ x_2x_{14} + x_4x_{16} = 0, \end{array} \right.$$

is equal to  $R_2 = 2^7 + 21 \cdot 2^5 + 490 \cdot 2^3 = 4720$

*Proof.* Set

$$\left\{ \begin{array}{l} Y_1 = x_1 + x_2 \cdot T \\ Y_2 = x_3 + x_4 \cdot T \\ Z_1 = x_5 + x_6 \cdot T \\ Z_2 = x_7 + x_8 \cdot T \\ U_1 = x_9 + x_{10} \cdot T \\ U_2 = x_{11} + x_{12} \cdot T \\ V_1 = x_{13} + x_{14} \cdot T \\ V_2 = x_{15} + x_{16} \cdot T \end{array} \right.$$

Then we obtain

$$\left\{ \begin{array}{l} Y_1Z_1 + Y_2Z_2 = 0, \\ Y_1U_1 + Y_2U_2 = 0, \\ Y_1V_1 + Y_2V_2 = 0 \end{array} \right. \Leftrightarrow \left\{ \begin{array}{l} x_1x_5 + x_3x_7 = 0, \\ x_1x_9 + x_3x_{11} = 0, \\ x_1x_{13} + x_3x_{15} = 0, \\ x_1x_6 + x_2x_5 + x_3x_8 + x_4x_7 = 0 \\ x_1x_{10} + x_2x_9 + x_3x_{12} + x_4x_{11} = 0 \\ x_1x_{14} + x_2x_{13} + x_3x_{16} + x_4x_{15} = 0 \\ x_2x_6 + x_4x_8 = 0, \\ x_2x_{10} + x_4x_{12} = 0, \\ x_2x_{14} + x_4x_{16} = 0, \end{array} \right.$$

□

*Example.* The number  $\Gamma_i \left[ \begin{smallmatrix} 2 & 2 \\ 2 & 2 \\ (3) \end{smallmatrix} \right] \times 4$  of rank  $i$  matrices of the form

$$\begin{pmatrix} \alpha_1 & \alpha_2 & \alpha_3 & \alpha_4 & \alpha_5 & \alpha_6 & \alpha_7 & \alpha_8 & \alpha_9 \\ \alpha_7 & \alpha_8 & \alpha_9 & \alpha_{10} & \alpha_{11} & \alpha_{12} & \alpha_{13} & \alpha_{14} & \alpha_{15} \\ \alpha_{13} & \alpha_{14} & \alpha_{15} & \alpha_{16} & \alpha_{17} & \alpha_{18} & \alpha_{19} & \alpha_{20} & \alpha_{21} \\ \alpha_{19} & \alpha_{20} & \alpha_{21} & \alpha_{22} & \alpha_{23} & \alpha_{24} & \alpha_{25} & \alpha_{26} & \alpha_{27} \end{pmatrix} \underset{\sim}{\text{rank}} \begin{pmatrix} \alpha_1 & \alpha_2 & \alpha_3 & \alpha_4 \\ \alpha_2 & \alpha_3 & \alpha_4 & \alpha_5 \\ \hline \beta_1 & \beta_2 & \beta_3 & \beta_4 \\ \beta_2 & \beta_3 & \beta_4 & \beta_5 \\ \hline \gamma_1 & \gamma_2 & \gamma_3 & \gamma_4 \\ \gamma_2 & \gamma_3 & \gamma_4 & \gamma_5 \\ \hline \delta_{11} & \delta_{12} & \delta_{13} & \delta_{14} \\ \delta_{21} & \delta_{22} & \delta_{23} & \delta_{24} \\ \delta_{31} & \delta_{32} & \delta_{33} & \delta_{34} \end{pmatrix}$$

is equal to

$$\begin{aligned} & 2^{3i} \cdot \Gamma_i \left[ \begin{smallmatrix} 2 \\ 2 \\ 2 \end{smallmatrix} \right] \times 4 \\ & + 7 \cdot 2^{(i-1)2} (2^4 - 2^{i-1}) \cdot \Gamma_{i-1} \left[ \begin{smallmatrix} 2 \\ 2 \\ 2 \end{smallmatrix} \right] \times 4 \\ & + 7 \cdot 2^{i-2} (2^4 - 2^{i-1}) (2^4 - 2^{i-2}) \cdot \Gamma_{i-2} \left[ \begin{smallmatrix} 2 \\ 2 \\ 2 \end{smallmatrix} \right] \times 4 \\ & + (2^4 - 2^{i-1}) (2^4 - 2^{i-2}) (2^4 - 2^{i-3}) \cdot \Gamma_{i-3} \left[ \begin{smallmatrix} 2 \\ 2 \\ 2 \end{smallmatrix} \right] \times 4 \quad \text{for } 0 \leq i \leq \inf(4, 9), \end{aligned}$$

Hence :

$$\Gamma_i \left[ \begin{smallmatrix} 2 & 2 \\ 2 & 2 \\ (3) \end{smallmatrix} \right] \times 4 = \begin{cases} 1 & \text{if } i = 0, \\ 273 & \text{if } i = 1, \\ 41062 & \text{if } i = 2, \\ 3807048 & \text{if } i = 3, \\ 130369344 & \text{if } i = 4. \end{cases}$$

For more details see [4]

## 5. THE INVERSE PROBLEM

How to compute the number  $R_q$  of solutions

$(Y_1, Z_1, U_1, V_1, \dots, Y_q, Z_q, U_q, V_q) \in (\mathbb{F}_2[T])^{4q}$  of the polynomial equations

$$\begin{cases} Y_1 Z_1 + Y_2 Z_2 + \dots + Y_q Z_q = 0, \\ Y_1 U_1 + Y_2 U_2 + \dots + Y_q U_q = 0, \\ Y_1 V_1 + Y_2 V_2 + \dots + Y_q V_q = 0, \end{cases}$$

satisfying the degree conditions

$$\deg Y_j \leq k-1, \quad \deg Z_j \leq s-1, \quad \deg U_j \leq s+m-1, \quad \deg V_j \leq s+m+l-1 \quad \text{for } 1 \leq j \leq q.$$

**Response:**

We need only to compute the number  $\Gamma_i^{\left[ \begin{smallmatrix} s \\ s+m \end{smallmatrix} \right]_{s+m+l}} \times k$  of triple persymmetric  $(3s+2m+l) \times k$  rank  $i$  matrices over  $\mathbb{F}_2$  for  $0 \leq i \leq \inf(k, 3s+2m+l)$  of the form

$$\left( \begin{array}{ccccc} \alpha_1 & \alpha_2 & \dots & \alpha_{k-1} & \alpha_k \\ \alpha_2 & \alpha_3 & \dots & \alpha_k & \alpha_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \alpha_{s-1} & \alpha_s & \dots & \alpha_{s+k-3} & \alpha_{s+k-2} \\ \alpha_s & \alpha_{s+1} & \dots & \alpha_{s+k-2} & \alpha_{s+k-1} \\ \hline \beta_1 & \beta_2 & \dots & \beta_{k-1} & \beta_k \\ \beta_2 & \beta_3 & \dots & \beta_k & \beta_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{m+1} & \beta_{m+2} & \dots & \beta_{k+m-1} & \beta_{k+m} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \beta_{s+m-1} & \beta_{s+m} & \dots & \beta_{s+m+k-3} & \beta_{s+m+k-2} \\ \beta_{s+m} & \beta_{s+m+1} & \dots & \beta_{s+m+k-2} & \beta_{s+m+k-1} \\ \hline \gamma_1 & \gamma_2 & \dots & \gamma_{k-1} & \gamma_k \\ \gamma_2 & \gamma_3 & \dots & \gamma_k & \gamma_{k+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \gamma_{m+1} & \gamma_{m+2} & \dots & \gamma_{k+m-1} & \gamma_{k+m} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \gamma_{s+m-1} & \gamma_{s+m} & \dots & \gamma_{s+m+k-3} & \gamma_{s+m+k-2} \\ \gamma_{s+m} & \gamma_{s+m+1} & \dots & \gamma_{s+m+k-2} & \gamma_{s+m+k-1} \\ \gamma_{s+m+1} & \gamma_{s+m+2} & \dots & \gamma_{s+m+k-1} & \gamma_{s+m+k} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \gamma_{s+m+l} & \gamma_{s+m+l+1} & \dots & \gamma_{s+m+l+k-2} & \gamma_{s+m+l+k-1} \end{array} \right).$$

To compute those numbers we use the following reductions formulas:

$$\begin{aligned}
\Gamma_{2s+1+m+j}^{\left[ \begin{smallmatrix} s \\ s+m+l \end{smallmatrix} \right] \times k} &= 16^j \cdot \Gamma_{2s+1+m}^{\left[ \begin{smallmatrix} s \\ s+m+l-j \end{smallmatrix} \right] \times (k-j)} && \text{if } 0 \leq j \leq l, k \geq 2s+1+m+j \\
\Gamma_{2s+1+m+l+j}^{\left[ \begin{smallmatrix} s \\ s+m+l \end{smallmatrix} \right] \times k} &= 16^{2j+l} \cdot \Gamma_{2s+1+m-j}^{\left[ \begin{smallmatrix} s \\ s+m-j \end{smallmatrix} \right] \times (k-2j-l)} && \text{if } 0 \leq j \leq m, k \geq 2s+1+m+l+j \\
\Gamma_{2s+1+2m+l+j}^{\left[ \begin{smallmatrix} s \\ s+m+l \end{smallmatrix} \right] \times k} &= 16^{2m+l+3j} \cdot \Gamma_{2(s-j)+1}^{\left[ \begin{smallmatrix} s-j \\ s-j \end{smallmatrix} \right] \times (k-2m-l-3j)} && \text{if } 0 \leq j \leq s-1, k \geq 2s+1+2m+l+j
\end{aligned}$$

Then  $R_q$  is equal to a linear combination of the  $\Gamma_i^{\left[ \begin{smallmatrix} s \\ s+m+l \end{smallmatrix} \right] \times k}$  for  $0 \leq i \leq \inf(k, 3s+2m+l)$   
More precisely:

$$R_q = 2^{(k+3s+2m+l)q - (3k+3s+2m+l-3)} \sum_{i=0}^{\inf(k, 3s+2m+l)} \Gamma_i^{\left[ \begin{smallmatrix} s \\ s+m+l \end{smallmatrix} \right] \times k} 2^{-iq}$$

**Remark :**

We have computed the  $\Gamma_i^{\left[ \begin{smallmatrix} s \\ s+m+l \end{smallmatrix} \right] \times k}$  for  $0 \leq i \leq \inf(k, 3s+2m+l)$  in the case  $l = 0$ . [see (4)]

**5.1. The inverse problem in the case  $s = 2, m = 3, l \geq 4$ .** The results in this subsection are new.

*Example.* To compute the number  $R_q$  of solutions  $(Y_1, Z_1, U_1, V_1, \dots, Y_q, Z_q, U_q, V_q) \in (\mathbb{F}_2[T])^{4q}$  of the polynomial equations

$$\begin{cases} Y_1 Z_1 + Y_2 Z_2 + \dots + Y_q Z_q = 0, \\ Y_1 U_1 + Y_2 U_2 + \dots + Y_q U_q = 0, \\ Y_1 V_1 + Y_2 V_2 + \dots + Y_q V_q = 0, \end{cases}$$

satisfying the degree conditions

$$\deg Y_j \leq k-1, \quad \deg Z_j \leq 1, \quad \deg U_j \leq 4, \quad \deg V_j \leq 4+l \quad \text{for } 1 \leq j \leq q,$$

we need only to compute the number  $\Gamma_i^{\left[ \begin{smallmatrix} 2 \\ 2+3+l \end{smallmatrix} \right] \times k}$  of triple persymmetric

$(12 + l) \times k$  rank  $i$  matrices over  $\mathbb{F}_2$  for  $0 \leq i \leq \inf(k, 12 + l)$  of the form

$$\begin{pmatrix} \alpha_1 & \alpha_2 & \dots & \alpha_{k-1} & \alpha_k \\ \alpha_2 & \alpha_3 & \dots & \alpha_k & \alpha_{k+1} \\ \hline \beta_1 & \beta_2 & \dots & \beta_{k-1} & \beta_k \\ \beta_2 & \beta_3 & \dots & \beta_k & \beta_{k+1} \\ \beta_3 & \beta_4 & \dots & \beta_{k+1} & \beta_{k+2} \\ \beta_4 & \beta_5 & \dots & \beta_{k+2} & \beta_{k+3} \\ \beta_5 & \beta_6 & \dots & \beta_{k+3} & \beta_{k+4} \\ \hline \gamma_1 & \gamma_2 & \dots & \gamma_{k-1} & \gamma_k \\ \gamma_2 & \gamma_3 & \dots & \gamma_k & \gamma_{k+1} \\ \gamma_3 & \gamma_4 & \dots & \gamma_{k+1} & \gamma_{k+2} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \gamma_{5+l} & \gamma_{6+l} & \dots & \gamma_{k+l+3} & \gamma_{k+l+4} \end{pmatrix}.$$

Proceeding as in (4), using the following reductions formulas:

$$\begin{aligned} \Gamma_{8+j}^{\left[ \begin{smallmatrix} 2 \\ 2+3 \\ 2+3+l \end{smallmatrix} \right] \times k} &= 16^j \cdot \Gamma_8^{\left[ \begin{smallmatrix} 2 \\ 2+3 \\ 2+3+l-j \end{smallmatrix} \right] \times (k-j)} && \text{if } 0 \leq j \leq l, k \geq 8 + j \\ \Gamma_{8+l+j}^{\left[ \begin{smallmatrix} 2 \\ 2+3 \\ 2+3+l \end{smallmatrix} \right] \times k} &= 16^{2j+l} \cdot \Gamma_{8-j}^{\left[ \begin{smallmatrix} 2 \\ 2+3-j \\ 2+3-j \end{smallmatrix} \right] \times (k-2j-l)} && \text{if } 0 \leq j \leq 3, k \geq 8 + l + j \\ \Gamma_{11+l+j}^{\left[ \begin{smallmatrix} 2 \\ 2+3 \\ 2+3+l \end{smallmatrix} \right] \times k} &= 16^{6+l+3j} \cdot \Gamma_{2(2-j)+1}^{\left[ \begin{smallmatrix} 2-j \\ 2-j \\ 2-j \end{smallmatrix} \right] \times (k-6-l-3j)} && \text{if } 0 \leq j \leq 1, k \geq 11 + l + j \end{aligned}$$

we obtain:

$$\begin{aligned} \Gamma_i^{\left[ \begin{smallmatrix} 2 \\ 2+3 \\ 2+3+l \end{smallmatrix} \right] \times k} &= \begin{cases} 1 & \text{if } i = 0, k > 0 \\ 21 & \text{if } i = 1, k > 1 \\ 2^{k+1} + r_2 (= 362) & \text{if } i = 2, k > 2 \\ 9 \cdot 2^{k+1} + r_3 (= 6048) & \text{if } i = 3, k > 3 \\ 39 \cdot 2^{k+2} + r_4 (= 98784) & \text{if } i = 4, k > 4 \\ 97 \cdot 2^{k+4} + r_5 (= 1580288) & \text{if } i = 5, k > 5 \\ 225 \cdot 2^{k+6} + r_6 (= 25135104) & \text{if } i = 6, k > 6 \\ 2^{2k+5} + 417 \cdot 2^{k+8} + r_7 (= 402571264 = 24571 \cdot 2^{14}) & \text{if } i = 7, k > 7 \end{cases} \\ \Gamma_{8+i}^{\left[ \begin{smallmatrix} 2 \\ 2+3 \\ 2+3+l \end{smallmatrix} \right] \times k} &= \begin{cases} 3 \cdot 2^{2k+5+2i} + 105 \cdot 2^{k+13+3i} + r_{8+i} (= 12285 \cdot 2^{19+4i}) & \text{if } 0 \leq i \leq l-4, k > 8+i \\ 3 \cdot 2^{2k+2l-1} + 137 \cdot 2^{k+3l+4} + r_{8+l-3} (= 12029 \cdot 2^{23+4 \cdot (l-4)}) & \text{if } i = l-3, k > 8+i \\ 3 \cdot 2^{2k+2l+1} + 213 \cdot 2^{k+3l+7} + r_{8+l-2} (= 11373 \cdot 2^{27+4 \cdot (l-4)}) & \text{if } i = l-2, k > 8+i \\ 11 \cdot 2^{2k+2l+3} + 333 \cdot 2^{k+3l+9} + r_{8+l-1} (= 10159 \cdot 2^{31+4 \cdot (l-4)}) & \text{if } i = l-1, k > 8+i \end{cases} \\ \Gamma_{8+l+i}^{\left[ \begin{smallmatrix} 2 \\ 2+3 \\ 2+3+l \end{smallmatrix} \right] \times k} &= \begin{cases} 21 \cdot 2^{2k+2l+5} + 2331 \cdot 2^{k+3l+11} + r_{8+l} (= 15435 \cdot 2^{34+4 \cdot (l-4)}) & \text{if } i = 0, k > 8+l+i \\ 21 \cdot 2^{2k+2l+8} + 2163 \cdot 2^{k+3l+15} + r_{9+l} (= 2835 \cdot 2^{39+4 \cdot (l-4)}) & \text{if } i = 1, k > 8+l+i \\ 53 \cdot 2^{2k+2l+11} + 1311 \cdot 2^{k+3l+19} + r_{10+l} (= -1417 \cdot 2^{44+4 \cdot (l-4)}) & \text{if } i = 2, k > 8+l+i \end{cases} \end{aligned}$$

$$\Gamma_{11+l+i}^{\left[ \begin{smallmatrix} 2 \\ 2+3 \\ 2+3+l \end{smallmatrix} \right] \times k} = \begin{cases} 105 \cdot 2^{2k+2l+14} - 315 \cdot 2^{k+3l+23} + r_{11+l} (= 105 \cdot 2^{49+4 \cdot (l-4)}) & \text{if } i = 0, k > 11 + l + i \\ 2^{3k+l+9} - 7 \cdot 2^{2k+2l+18} + 7 \cdot 2^{k+3l+28} + r_{12+l} (= -2^{55+4 \cdot (l-4)}) & \text{if } i = 1, k \geq 11 + l + i \end{cases}$$

Then we get:

$$R_q = 2^{(k+12+l)q-(3k+l+9)} \sum_{i=0}^{\inf(k,12+l)} \Gamma_i^{\left[ \begin{smallmatrix} 2 \\ 2+3 \\ 2+3+l \end{smallmatrix} \right] \times k} 2^{-iq}$$

**5.2. The inverse problem in the case  $s = 2$ ,  $m = 3$ ,  $l = 4$ ,  $k = 6$ .** The results in this subsection are new.

*Example.* We get:

$$\Gamma_i^{\left[ \begin{smallmatrix} 2 \\ 2+3 \\ 2+3+4 \end{smallmatrix} \right] \times 6} = \begin{cases} 1 & \text{if } i = 0, \\ 21 & \text{if } i = 1, \\ 490 & \text{if } i = 2, \\ 7200 & \text{if } i = 3, \\ 108768 & \text{if } i = 4, \\ 1679616 & \text{if } i = 5, \\ 2145687552 & \text{if } i = 6, \end{cases}$$

The number  $R_q$  of solutions  $(Y_1, Z_1, U_1, V_1, \dots, Y_q, Z_q, U_q, V_q) \in (\mathbb{F}_2[T])^{4q}$  of the polynomial equations

$$\begin{cases} Y_1 Z_1 + Y_2 Z_2 + \dots + Y_q Z_q = 0, \\ Y_1 U_1 + Y_2 U_2 + \dots + Y_q U_q = 0, \\ Y_1 V_1 + Y_2 V_2 + \dots + Y_q V_q = 0, \end{cases}$$

satisfying the degree conditions

$$\deg Y_i \leq 5, \quad \deg Z_i \leq 1, \quad \deg U_i \leq 4, \quad \deg V_i \leq 8 \quad \text{for } 1 \leq i \leq q.$$

is equal to

$$2^{22q-31} \sum_{i=0}^6 \Gamma_i^{\left[ \begin{smallmatrix} 2 \\ 2+3 \\ 2+3+4 \end{smallmatrix} \right] \times 6} 2^{-iq}$$



**5.3. The inverse problem in the case  $s = 3$ ,  $m = 0$ ,  $l \geq 0$ .** The results in this subsection are new.

*Example.*  $s = 3$ ,  $m = 0$ ,  $l = 0$ .

The number  $\Gamma_i^{\left[\begin{smallmatrix} 3 \\ 3 \\ 3 \end{smallmatrix}\right] \times k}$  of triple persymmetric  $9 \times k$  rank  $i$  matrices over  $\mathbb{F}_2$  of the form

$$\begin{pmatrix} \alpha_1 & \alpha_2 & \dots & \dots & \alpha_{k-1} & \alpha_k \\ \alpha_2 & \alpha_3 & \dots & \dots & \alpha_k & \alpha_{k+1} \\ \alpha_3 & \alpha_4 & \dots & \dots & \alpha_{k+1} & \alpha_{k+2} \\ \hline \beta_1 & \beta_2 & \dots & \dots & \beta_{k-1} & \beta_k \\ \beta_2 & \beta_3 & \dots & \dots & \beta_k & \beta_{k+1} \\ \beta_3 & \beta_4 & \dots & \dots & \beta_{k+1} & \beta_{k+2} \\ \hline \gamma_1 & \gamma_2 & \dots & \dots & \gamma_{k-1} & \gamma_k \\ \gamma_2 & \gamma_3 & \dots & \dots & \gamma_k & \gamma_{k+1} \\ \gamma_3 & \gamma_4 & \dots & \dots & \gamma_{k+1} & \gamma_{k+2} \end{pmatrix}.$$

is equal to

$$\Gamma_i^{\left[\begin{smallmatrix} 3 \\ 3 \\ 3 \end{smallmatrix}\right] \times k} = \begin{cases} 1 & \text{if } i = 0 \\ 21 & \text{if } i = 1 \\ 378 & \text{if } i = 2 \\ 7 \cdot 2^{k+2} + 5936 & \text{if } i = 3 \\ 147 \cdot 2^{k+2} + 84672 & \text{if } i = 4 \\ 147 \cdot 9 \cdot 2^{k+3} + 959616 & \text{if } i = 5 \\ 7 \cdot 2^{2k+4} + 2121 \cdot 2^{k+6} + 5863424 & \text{if } i = 6 \\ 105 \cdot 2^{2k+4} + 2625 \cdot 2^{k+9} - 92897280 & \text{if } i = 7 \\ 105 \cdot 2^{2k+8} - 315 \cdot 2^{k+14} + 220200960 & \text{if } i = 8 \\ 2^{3k+6} - 7 \cdot 2^{2k+12} + 7 \cdot 2^{k+19} - 134217728 & \text{if } i = 9, k \geq 9 \end{cases}$$

*Example.*  $s = 3$ ,  $m = 0$ ,  $l = 1$ .

The number  $\Gamma_i^{\left[\begin{smallmatrix} 3 \\ 3 \\ 3 \\ 3+1 \end{smallmatrix}\right] \times k}$  of triple persymmetric  $10 \times k$  rank  $i$  matrices over  $\mathbb{F}_2$  of the form

$$\begin{pmatrix} \alpha_1 & \alpha_2 & \dots & \dots & \alpha_{k-1} & \alpha_k \\ \alpha_2 & \alpha_3 & \dots & \dots & \alpha_k & \alpha_{k+1} \\ \alpha_3 & \alpha_4 & \dots & \dots & \alpha_{k+1} & \alpha_{k+2} \\ \hline \beta_1 & \beta_2 & \dots & \dots & \beta_{k-1} & \beta_k \\ \beta_2 & \beta_3 & \dots & \dots & \beta_k & \beta_{k+1} \\ \beta_3 & \beta_4 & \dots & \dots & \beta_{k+1} & \beta_{k+2} \\ \hline \gamma_1 & \gamma_2 & \dots & \dots & \gamma_{k-1} & \gamma_k \\ \gamma_2 & \gamma_3 & \dots & \dots & \gamma_k & \gamma_{k+1} \\ \gamma_3 & \gamma_4 & \dots & \dots & \gamma_{k+1} & \gamma_{k+2} \\ \gamma_4 & \gamma_5 & \dots & \dots & \gamma_{k+2} & \gamma_{k+3} \end{pmatrix}.$$

is equal to

$$\Gamma_i^{\left[ \begin{smallmatrix} 3 \\ 3 \\ 3+1 \end{smallmatrix} \right] \times k} = \begin{cases} 1 & \text{if } i = 0, \\ 21 & \text{if } i = 1, \\ 378 & \text{if } i = 2, \\ 3 \cdot 2^{k+2} + 6192 & \text{if } i = 3, \\ 71 \cdot 2^{k+2} + 124864 & \text{if } i = 4, \\ 651 \cdot 2^{k+3} + 1246848 & \text{if } i = 5, \\ 2^{2k+4} + 645 \cdot 2^{k+7} + 15464448 & \text{if } i = 6, \\ 27 \cdot 2^{2k+4} + 531 \cdot 2^{k+11} + 93782016 & \text{if } i = 7, \\ 105 \cdot 2^{2k+6} + 2625 \cdot 2^{k+12} - 1486356480 & \text{if } i = 8, \\ 105 \cdot 2^{2k+10} - 315 \cdot 2^{k+17} + 3523215360 & \text{if } i = 9, \\ 2^{3k+7} - 7 \cdot 2^{2k+14} + 7 \cdot 2^{k+22} - 2147483648 & \text{if } i = 10 \end{cases}$$

*Example.*  $s = 3$ ,  $m = 0$ ,  $l = 2$ .

The number  $\Gamma_i^{\left[ \begin{smallmatrix} 3 \\ 3 \\ 3+2 \end{smallmatrix} \right] \times k}$  of triple persymmetric  $11 \times k$  rank  $i$  matrices over  $\mathbb{F}_2$  of the form

$$\begin{pmatrix} \alpha_1 & \alpha_2 & \dots & \dots & \alpha_{k-1} & \alpha_k \\ \alpha_2 & \alpha_3 & \dots & \dots & \alpha_k & \alpha_{k+1} \\ \alpha_3 & \alpha_4 & \dots & \dots & \alpha_{k+1} & \alpha_{k+2} \\ \hline \beta_1 & \beta_2 & \dots & \dots & \beta_{k-1} & \beta_k \\ \beta_2 & \beta_3 & \dots & \dots & \beta_k & \beta_{k+1} \\ \beta_3 & \beta_4 & \dots & \dots & \beta_{k+1} & \beta_{k+2} \\ \hline \gamma_1 & \gamma_2 & \dots & \dots & \gamma_{k-1} & \gamma_k \\ \gamma_2 & \gamma_3 & \dots & \dots & \gamma_k & \gamma_{k+1} \\ \gamma_3 & \gamma_4 & \dots & \dots & \gamma_{k+1} & \gamma_{k+2} \\ \gamma_4 & \gamma_5 & \dots & \dots & \gamma_{k+2} & \gamma_{k+3} \\ \gamma_5 & \gamma_6 & \dots & \dots & \gamma_{k+3} & \gamma_{k+4} \end{pmatrix}.$$

is equal to

$$\Gamma_i^{\left[ \begin{smallmatrix} 3 \\ 3 \\ 3+2 \end{smallmatrix} \right] \times k} = \begin{cases} 1 & \text{if } i = 0, \\ 21 & \text{if } i = 1, \\ 378 & \text{if } i = 2, \\ 3 \cdot 2^{k+2} + 6192 & \text{if } i = 3, \\ 39 \cdot 2^{k+2} + 99264 & \text{if } i = 4, \\ 347 \cdot 2^{k+3} + 1503872 & \text{if } i = 5, \\ 2^{2k+4} + 618 \cdot 2^{k+6} + 21426176 & \text{if } i = 6, \\ 3 \cdot 2^{2k+4} + 1293 \cdot 2^{k+9} + 246448128 & \text{if } i = 7, \\ 27 \cdot 2^{2k+6} + 531 \cdot 2^{k+14} + 1500512256 & \text{if } i = 8, \\ 105 \cdot 2^{2k+8} + 2625 \cdot 2^{k+15} - 92897280 \cdot 2^8 & \text{if } i = 9, \\ 105 \cdot 2^{2k+12} - 315 \cdot 2^{k+20} + 53760 \cdot 2^{20} & \text{if } i = 10, \\ 2^{3k+8} - 7 \cdot 2^{2k+16} + 7 \cdot 2^{k+25} - 2^{35} & \text{if } i = 11 \end{cases}$$

*Example.*  $s = 3$ ,  $m = 0$ ,  $l = 3$ .

The number  $\Gamma_i^{\left[ \begin{smallmatrix} 3 \\ 3 \\ 3+3 \end{smallmatrix} \right] \times k}$  of triple persymmetric  $12 \times k$  rank  $i$  matrices over  $\mathbb{F}_2$  of the form

$$\begin{pmatrix} \alpha_1 & \alpha_2 & \dots & \dots & \alpha_{k-1} & \alpha_k \\ \alpha_2 & \alpha_3 & \dots & \dots & \alpha_k & \alpha_{k+1} \\ \alpha_3 & \alpha_4 & \dots & \dots & \alpha_{k+1} & \alpha_{k+2} \\ \hline \beta_1 & \beta_2 & \dots & \dots & \beta_{k-1} & \beta_k \\ \beta_2 & \beta_3 & \dots & \dots & \beta_k & \beta_{k+1} \\ \beta_3 & \beta_4 & \dots & \dots & \beta_{k+1} & \beta_{k+2} \\ \hline \gamma_1 & \gamma_2 & \dots & \dots & \gamma_{k-1} & \gamma_k \\ \gamma_2 & \gamma_3 & \dots & \dots & \gamma_k & \gamma_{k+1} \\ \gamma_3 & \gamma_4 & \dots & \dots & \gamma_{k+1} & \gamma_{k+2} \\ \gamma_4 & \gamma_5 & \dots & \dots & \gamma_{k+2} & \gamma_{k+3} \\ \gamma_5 & \gamma_6 & \dots & \dots & \gamma_{k+3} & \gamma_{k+4} \\ \gamma_6 & \gamma_7 & \dots & \dots & \gamma_{k+4} & \gamma_{k+5} \end{pmatrix}.$$

is equal to

$$\Gamma_i^{\left[ \begin{smallmatrix} 3 & 3 & 3 \\ 3+3 \end{smallmatrix} \right] \times k} = \begin{cases} 1 & \text{if } i = 0, \\ 21 & \text{if } i = 1, \\ 378 & \text{if } i = 2, \\ 3 \cdot 2^{k+2} + 6192 & \text{if } i = 3, \\ 39 \cdot 2^{k+2} + 99264 & \text{if } i = 4, \\ 219 \cdot 2^{k+3} + 1569408 & \text{if } i = 5, \\ 2^{2k+4} + 314 \cdot 2^{k+6} + 24113152 & \text{if } i = 6, \\ 3 \cdot 2^{2k+4} + 621 \cdot 2^{k+9} + 342786048 & \text{if } i = 7, \\ 3 \cdot 2^{2k+6} + 1293 \cdot 2^{k+12} + 3943170048 & \text{if } i = 8, \\ 27 \cdot 2^{2k+8} + 531 \cdot 2^{k+17} + 93782016 \cdot 2^8 & \text{if } i = 9, \\ 105 \cdot 2^{2k+10} + 2625 \cdot 2^{k+18} - 92897280 \cdot 2^{12} & \text{if } i = 10, \\ 105 \cdot 2^{2k+14} - 315 \cdot 2^{k+23} + 53760 \cdot 2^{24} & \text{if } i = 11, \\ 2^{3k+9} - 7 \cdot 2^{2k+18} + 7 \cdot 2^{k+28} - 2^{39} & \text{if } i = 12 \end{cases}$$

*Example.*  $s = 3$ ,  $m = 0$ ,  $l = 4$ .

The number  $\Gamma_i^{\left[ \begin{smallmatrix} 3 & 3 & 3 \\ 3+4 \end{smallmatrix} \right] \times k}$  of triple persymmetric  $13 \times k$  rank  $i$  matrices over  $\mathbb{F}_2$  of the form

$$\begin{pmatrix} \alpha_1 & \alpha_2 & \dots & \dots & \alpha_{k-1} & \alpha_k \\ \alpha_2 & \alpha_3 & \dots & \dots & \alpha_k & \alpha_{k+1} \\ \alpha_3 & \alpha_4 & \dots & \dots & \alpha_{k+1} & \alpha_{k+2} \\ \hline \beta_1 & \beta_2 & \dots & \dots & \beta_{k-1} & \beta_k \\ \beta_2 & \beta_3 & \dots & \dots & \beta_k & \beta_{k+1} \\ \beta_3 & \beta_4 & \dots & \dots & \beta_{k+1} & \beta_{k+2} \\ \hline \gamma_1 & \gamma_2 & \dots & \dots & \gamma_{k-1} & \gamma_k \\ \gamma_2 & \gamma_3 & \dots & \dots & \gamma_k & \gamma_{k+1} \\ \gamma_3 & \gamma_4 & \dots & \dots & \gamma_{k+1} & \gamma_{k+2} \\ \gamma_4 & \gamma_5 & \dots & \dots & \gamma_{k+2} & \gamma_{k+3} \\ \gamma_5 & \gamma_6 & \dots & \dots & \gamma_{k+3} & \gamma_{k+4} \\ \gamma_6 & \gamma_7 & \dots & \dots & \gamma_{k+4} & \gamma_{k+5} \\ \gamma_7 & \gamma_8 & \dots & \dots & \gamma_{k+5} & \gamma_{k+6} \end{pmatrix}.$$

is equal to

$$\Gamma_i^{\left[ \begin{smallmatrix} 3 \\ 3 \\ 3 \end{smallmatrix} \right]_{3+4} \times k} = \begin{cases} 1 & \text{if } i = 0, \\ 21 & \text{if } i = 1, \\ 378 & \text{if } i = 2, \\ 3 \cdot 2^{k+2} + 6192 & \text{if } i = 3, \\ 39 \cdot 2^{k+2} + 99264 & \text{if } i = 4, \\ 219 \cdot 2^{k+3} + 1569408 & \text{if } i = 5, \\ 2^{2k+4} + 186 \cdot 2^{k+6} + 25161728 & \text{if } i = 6, \\ 3 \cdot 2^{2k+4} + 317 \cdot 2^{k+9} + 38577664 & \text{if } i = 7, \\ 3 \cdot 2^{2k+6} + 621 \cdot 2^{k+12} + 5356032 \cdot 2^{10} & \text{if } i = 8, \\ 3 \cdot 2^{2k+8} + 1293 \cdot 2^{k+15} + 240672 \cdot 2^{18} & \text{if } i = 9, \\ 27 \cdot 2^{2k+10} + 531 \cdot 2^{k+20} + 93782016 \cdot 2^{12} & \text{if } i = 10, \\ 105 \cdot 2^{2k+12} + 2625 \cdot 2^{k+21} - 92897280 \cdot 2^{16} & \text{if } i = 11, \\ 105 \cdot 2^{2k+16} - 315 \cdot 2^{k+26} + 53760 \cdot 2^{28} & \text{if } i = 12, \\ 2^{3k+10} - 7 \cdot 2^{2k+20} + 7 \cdot 2^{k+31} - 2^{43} & \text{if } i = 13 \end{cases}$$

*Example.*  $s = 3$ ,  $m = 0$ ,  $l = 5$ .

The number  $\Gamma_i^{\left[ \begin{smallmatrix} 3 \\ 3 \\ 3 \end{smallmatrix} \right]_{3+5} \times k}$  of triple persymmetric  $14 \times k$  rank  $i$  matrices over  $\mathbb{F}_2$  of the form

$$\begin{pmatrix} \alpha_1 & \alpha_2 & \dots & \dots & \alpha_{k-1} & \alpha_k \\ \alpha_2 & \alpha_3 & \dots & \dots & \alpha_k & \alpha_{k+1} \\ \alpha_3 & \alpha_4 & \dots & \dots & \alpha_{k+1} & \alpha_{k+2} \\ \hline \beta_1 & \beta_2 & \dots & \dots & \beta_{k-1} & \beta_k \\ \beta_2 & \beta_3 & \dots & \dots & \beta_k & \beta_{k+1} \\ \beta_3 & \beta_4 & \dots & \dots & \beta_{k+1} & \beta_{k+2} \\ \hline \gamma_1 & \gamma_2 & \dots & \dots & \gamma_{k-1} & \gamma_k \\ \gamma_2 & \gamma_3 & \dots & \dots & \gamma_k & \gamma_{k+1} \\ \gamma_3 & \gamma_4 & \dots & \dots & \gamma_{k+1} & \gamma_{k+2} \\ \gamma_4 & \gamma_5 & \dots & \dots & \gamma_{k+2} & \gamma_{k+3} \\ \gamma_5 & \gamma_6 & \dots & \dots & \gamma_{k+3} & \gamma_{k+4} \\ \gamma_6 & \gamma_7 & \dots & \dots & \gamma_{k+4} & \gamma_{k+5} \\ \gamma_7 & \gamma_8 & \dots & \dots & \gamma_{k+5} & \gamma_{k+6} \\ \gamma_8 & \gamma_9 & \dots & \dots & \gamma_{k+6} & \gamma_{k+7} \end{pmatrix}.$$

is equal to

$$\Gamma_i^{\left[ \begin{smallmatrix} 3 \\ 3 \\ 3 \end{smallmatrix} \right]_{3+5} \times k} = \begin{cases} 1 & \text{if } i = 0, \\ 21 & \text{if } i = 1, \\ 378 & \text{if } i = 2, \\ 3 \cdot 2^{k+2} + 6192 & \text{if } i = 3, \\ 39 \cdot 2^{k+2} + 99264 & \text{if } i = 4, \\ 219 \cdot 2^{k+3} + 1569408 & \text{if } i = 5, \\ 2^{2k+4} + 186 \cdot 2^{k+6} + 25161728 & \text{if } i = 6, \\ 3 \cdot 2^{2k+4} + 189 \cdot 2^{k+9} + 402554880 & \text{if } i = 7, \\ 3 \cdot 2^{2k+6} + 317 \cdot 2^{k+12} + 385777664 \cdot 2^4 & \text{if } i = 8, \\ 3 \cdot 2^{2k+8} + 621 \cdot 2^{k+15} + 342786048 \cdot 2^8 & \text{if } i = 9, \\ 3 \cdot 2^{2k+10} + 1293 \cdot 2^{k+18} + 246448128 \cdot 2^{12} & \text{if } i = 10, \\ 27 \cdot 2^{2k+12} + 531 \cdot 2^{k+23} + 93782016 \cdot 2^{16} & \text{if } i = 11, \\ 105 \cdot 2^{2k+14} + 2625 \cdot 2^{k+24} - 92897280 \cdot 2^{20} & \text{if } i = 12, \\ 105 \cdot 2^{2k+18} - 315 \cdot 2^{k+29} + 53760 \cdot 2^{32} & \text{if } i = 13, \\ 2^{3k+11} - 7 \cdot 2^{2k+22} + 7 \cdot 2^{k+34} - 2^{47} & \text{if } i = 14 \end{cases}$$

*Example.*  $s = 3$ ,  $m = 0$ ,  $l \geq 5$

The number  $\Gamma_i^{\left[ \begin{smallmatrix} 3 \\ 3 \\ 3 \end{smallmatrix} \right]_{3+l} \times k}$  of triple persymmetric  $(9 + l) \times k$  rank  $i$  matrices over  $\mathbb{F}_2$  of the form

$$\begin{pmatrix} \alpha_1 & \alpha_2 & \dots & \dots & \alpha_{k-1} & \alpha_k \\ \alpha_2 & \alpha_3 & \dots & \dots & \alpha_k & \alpha_{k+1} \\ \alpha_3 & \alpha_4 & \dots & \dots & \alpha_{k+1} & \alpha_{k+2} \\ \hline \beta_1 & \beta_2 & \dots & \dots & \beta_{k-1} & \beta_k \\ \beta_2 & \beta_3 & \dots & \dots & \beta_k & \beta_{k+1} \\ \beta_3 & \beta_4 & \dots & \dots & \beta_{k+1} & \beta_{k+2} \\ \hline \gamma_1 & \gamma_2 & \dots & \dots & \gamma_{k-1} & \gamma_k \\ \gamma_2 & \gamma_3 & \dots & \dots & \gamma_k & \gamma_{k+1} \\ \gamma_3 & \gamma_4 & \dots & \dots & \gamma_{k+1} & \gamma_{k+2} \\ \gamma_4 & \gamma_5 & \dots & \dots & \gamma_{k+2} & \gamma_{k+3} \\ \gamma_5 & \gamma_6 & \dots & \dots & \gamma_{k+3} & \gamma_{k+4} \\ \gamma_6 & \gamma_7 & \dots & \dots & \gamma_{k+4} & \gamma_{k+5} \\ \gamma_7 & \gamma_8 & \dots & \dots & \gamma_{k+5} & \gamma_{k+6} \\ \gamma_8 & \gamma_9 & \dots & \dots & \gamma_{k+6} & \gamma_{k+7} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \gamma_{3+l} & \gamma_{4+l} & \dots & \dots & \gamma_{k+1+l} & \gamma_{k+2+l} \end{pmatrix}.$$

is equal to

$$\Gamma_i \begin{bmatrix} 3 \\ 3 \\ 3+l \end{bmatrix}^{\times k} = \begin{cases} 1 & \text{if } i = 0, \\ 21 & \text{if } i = 1, \\ 378 & \text{if } i = 2, \\ 3 \cdot 2^{k+2} + 6192 & \text{if } i = 3, \\ 39 \cdot 2^{k+2} + 99264 & \text{if } i = 4, \\ 219 \cdot 2^{k+3} + 1569408 & \text{if } i = 5, \\ 2^{2k+4} + 186 \cdot 2^{k+6} + 25161728 & \text{if } i = 6, \\ 3 \cdot 2^{2k+2i-10} + 189 \cdot 2^{k+3i-12} + 402554880 \cdot 2^{4(i-7)} & \text{if } 7 \leq i \leq l+2 \\ 3 \cdot 2^{2k+2l-4} + 317 \cdot 2^{k+3l-3} + 38577664 \cdot 2^{4l-16} & \text{if } i = l+3, \\ 3 \cdot 2^{2k+2l-2} + 621 \cdot 2^{k+3l} + 342786048 \cdot 2^{4l-12} & \text{if } i = l+4, \\ 3 \cdot 2^{2k+2l} + 1293 \cdot 2^{k+3l+3} + 246448128 \cdot 2^{4l-8} & \text{if } i = l+5, \\ 27 \cdot 2^{2k+2l+2} + 531 \cdot 2^{k+3l+8} + 93782016 \cdot 2^{4l-4} & \text{if } i = l+6, \\ 105 \cdot 2^{2k+2l+4} + 2625 \cdot 2^{k+3l+9} - 92897280 \cdot 2^{4l} & \text{if } i = l+7, \\ 105 \cdot 2^{2k+2l+8} - 315 \cdot 2^{k+3l+14} + 53760 \cdot 2^{4l+12} & \text{if } i = l+8 \\ 2^{3k+l+6} - 7 \cdot 2^{2k+2l+12} + 7 \cdot 2^{k+3l+19} - 2^{4l+27} & \text{if } i = l+9 \end{cases}$$

*Proof.* We compute  $\Gamma_i \begin{bmatrix} 3 \\ 3 \\ 3+l \end{bmatrix}^{\times k}$  for  $0 \leq i \leq l+9$  by using the following two reduction formulas:

$$\begin{aligned} \Gamma_{7+j} \begin{bmatrix} 3 \\ 3 \\ 3+l \end{bmatrix}^{\times k} &= 2^{4j} \cdot \Gamma_7 \begin{bmatrix} 3 \\ 3 \\ 3+l-j \end{bmatrix}^{\times (k-j)} && \text{if } 0 \leq j \leq l, k \geq 7+j \\ \Gamma_{7+l+j} \begin{bmatrix} 3 \\ 3 \\ 3+l \end{bmatrix}^{\times k} &= 2^{4l+12j} \cdot \Gamma_{2(3-j)+1} \begin{bmatrix} 3-j \\ 3-j \\ 3-j \end{bmatrix}^{\times (k-l-3j)} && \text{if } 0 \leq j \leq 2, k \geq 7+l+j \end{aligned}$$

combined with the formula:

$$\Gamma_7 \begin{bmatrix} 3 \\ 3 \\ 3+l \end{bmatrix}^{\times k} = \Gamma_7 \begin{bmatrix} 3 \\ 3 \\ 3+5 \end{bmatrix}^{\times k} \quad \text{if } l \geq 5, k > 7$$

More precisely : (with  $k > 7+j$ )

$$\Gamma_{7+j}^{\left[\begin{smallmatrix} 3 \\ 3+l \end{smallmatrix}\right] \times k} = \begin{cases} 2^{4j} \cdot \Gamma_7^{\left[\begin{smallmatrix} 3 \\ 3+5 \end{smallmatrix}\right] \times (k-j)} = 3 \cdot 2^{2k+2j+4} + 189 \cdot 2^{k+3j+9} + 402554880 \cdot 2^{4j} & \text{if } 0 \leq j \leq l-5, \\ 2^{4l-16} \cdot \Gamma_7^{\left[\begin{smallmatrix} 3 \\ 3+4 \end{smallmatrix}\right] \times (k-l+4)} = 3 \cdot 2^{2k+2l-4} + 317 \cdot 2^{k+3l-3} + 38577664 \cdot 2^{4l-16} & \text{if } j = l-4, \\ 2^{4l-12} \cdot \Gamma_7^{\left[\begin{smallmatrix} 3 \\ 3+3 \end{smallmatrix}\right] \times (k-l+3)} = 3 \cdot 2^{2k+2l-2} + 621 \cdot 2^{k+3l} + 342786048 \cdot 2^{4l-12} & \text{if } j = l-3, \\ 2^{4l-8} \cdot \Gamma_7^{\left[\begin{smallmatrix} 3 \\ 3+2 \end{smallmatrix}\right] \times (k-l+2)} = 3 \cdot 2^{2k+2l} + 1293 \cdot 2^{k+3l+3} + 246448128 \cdot 2^{4l-8} & \text{if } j = l-2, \\ 2^{4l-4} \cdot \Gamma_7^{\left[\begin{smallmatrix} 3 \\ 3+1 \end{smallmatrix}\right] \times (k-l+1)} = 27 \cdot 2^{2k+2l+2} + 531 \cdot 2^{k+3l+8} + 93782016 \cdot 2^{4l-4} & \text{if } j = l-1, \\ 2^{4l} \cdot \Gamma_7^{\left[\begin{smallmatrix} 3 \\ 3 \end{smallmatrix}\right] \times (k-l)} = 105 \cdot 2^{2k+2l+4} + 2625 \cdot 2^{k+3l+9} - 92897280 \cdot 2^{4l} & \text{if } j = l, \\ 2^{4l+12} \cdot \Gamma_5^{\left[\begin{smallmatrix} 2 \\ 2 \end{smallmatrix}\right] \times (k-l-3)} = 105 \cdot 2^{2k+2l+8} - 315 \cdot 2^{k+3l+14} + 53760 \cdot 2^{4l+12} & \text{if } j = l+1, \\ 2^{4l+24} \cdot \Gamma_3^{\left[\begin{smallmatrix} 1 \\ 1 \end{smallmatrix}\right] \times (k-l-6)} = 2^{3k+l+6} - 7 \cdot 2^{2k+2l+12} + 7 \cdot 2^{k+3l+19} - 2^{4l+27} & \text{if } j = l+2 \end{cases}$$

□

The number  $R_q$  of solutions

$(Y_1, Z_1, U_1, V_1, \dots, Y_q, Z_q, U_q, V_q) \in (\mathbb{F}_2[T])^{4q}$  of the polynomial equations

$$\begin{cases} Y_1 Z_1 + Y_2 Z_2 + \dots + Y_q Z_q = 0, \\ Y_1 U_1 + Y_2 U_2 + \dots + Y_q U_q = 0, \\ Y_1 V_1 + Y_2 V_2 + \dots + Y_q V_q = 0, \end{cases}$$

satisfying the degree conditions

$$\deg Y_j \leq k-1, \quad \deg Z_j \leq 2, \quad \deg U_j \leq 2, \quad \deg V_j \leq 2+l \quad \text{for } 1 \leq j \leq q,$$

is equal to

$$2^{(k+9+l)q-(3k+l+6)} \sum_{i=0}^{\inf(k,9+l)} \Gamma_i^{\left[\begin{smallmatrix} 3 \\ 3+l \end{smallmatrix}\right] \times k} 2^{-iq}$$

In particular:

$$R_1 = 2^{-2k+3} \sum_{i=0}^{\inf(k,9+l)} \Gamma_i^{\left[\begin{smallmatrix} 3 \\ 3+l \end{smallmatrix}\right] \times k} 2^{-i} = 2^{9+l} + 2^k - 1$$

## REFERENCES

- [1] Daykin David E, Distribution of Bordered Persymmetric Matrices in a finite field J. reine angew. Math, **203**(1960) ,47-54



- [2] Cherly, Jorgen.  
Exponential sums and rank of persymmetric matrices over  $\mathbf{F}_2$   
arXiv : 0711.1306, 46 pp
- [3] Cherly, Jorgen.  
Exponential sums and rank of double persymmetric matrices over  $\mathbf{F}_2$   
arXiv : 0711.1937, 160 pp
- [4] Cherly, Jorgen.  
Exponential sums and rank of triple persymmetric matrices over  $\mathbf{F}_2$   
arXiv : 0803.1398, 233 pp
- [5] Cherly, Jorgen.  
Results about persymmetric matrices over  $\mathbf{F}_2$  and related exponentials sums  
arXiv : 0803.2412v2, 32 pp

DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ DE BREST, 29238 BREST CEDEX 3,  
FRANCE

*E-mail address:* Jorgen.Cherly@univ-brest.fr

*E-mail address:* andersen69@wanadoo.fr